



| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|---------------------------|--|--|--|----------------|----|--------|
| CALCIUM SUPPLEMENT | | | | | | |
| 1 | BOCA Calcium emulsion "LONG LIFE" | Tricalcium phosphate 103mg, 1600 I.U. of Vitamin A and 160 I.U. of Vitamin D3/ml | For deficiency of calcium, Vit.A and Vit.D | Prescription | | |
| 2 | Concril Capsules 250mg | Crystalline D-Glucosamine Sulfate | Relieve the pain of degenerative arthritis. | OTC | | |
| PENICILLINS | | | | | | |
| 1 | AMPICILLIN SODIUM INJECTION 1G | AMPICILLIN SODIUM 1GM | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, Meningococcal and other susceptible bacteria | Prescription | | |
| 2 | AMOX INJECTION 500MG | AMOXICILLIN ANHYDROUS 500MG | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, Meningococcal and other susceptible bacteria. | Prescription | | |
| 3 | AMBACILLIN POWDER FOR INJECTION | AMPICILLIN 1GM/2GM + SULBACTAM 0.5GM/1GM | Infections caused by Streptococcus, Staphylococcus, Escherichia coli, Klebsiella, Serratia, Haemophilus influenzae, Proteus vulgaris, Citrobacter, Bacterioides Fusobacterium, Clostridium Difficile, Branhamella Catarrhalis. | Prescription | | |
| 4 | BETAMYCIN POWDER FOR INJECTION | PIPERACILLIN 2GM/3GM/4GM + TAZOBACTAM 0.25GM/0.375GM/0.5GM | The treatment of patients with moderate to severe infections caused by β -lactamase-producing strains who are susceptible to 1. piperacillin 2.piperacillin/tazobactam, but has resistance to piperacillin | Prescription | | |
| 5 | CO-TAZO POWDER FOR INJECTION | PIPERACILLIN 2GM/3GM/4GM + TAZOBACTAM 0.25GM/0.375GM/0.5GM | The treatment of patients with moderate to severe infections caused by β -lactamase-producing strains who are susceptible to 1. piperacillin 2.piperacillin/tazobactam, but has resistance to piperacillin | Prescription | | |
| 6 | Amonado for Intravenous Injection "L.L." | Amoxicillin 1g +Clavulanic Acid 0.2g | Infections caused by staphylococci, streptococci, pneumococci, meningococci and other susceptible bacteria. | Prescription | | |
| CEPHALOSPORINS | | | | | | |
| 1 | SEAFAR POWDER 0.5G FOR INJECTION | CEFAZOLIN (SODIUM) 0.5GM | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, meningococcal and other susceptible bacteria | Prescription | | |
| 2 | SEAFAR POWDER 1G/2G/5G/10G/20G FOR INJECTION | CEFAZOLIN (SODIUM) 1G/2G/5G/10G/20G | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, meningococcal and other susceptible bacteria | Prescription | | |
| 3 | CEDO INJECTION 1GM | CEFAMANDOLE 1G | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, Bacillus and other susceptible bacteria. | Prescription | | |
| 4 | CEFOTIN POWDER FOR INJECTION "CEFOXITIN" | CEFOXITIN 1G/2G | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, meningococcal and other susceptible bacteria. | Prescription | | |
| 5 | CEFURO INJECTION 250MG | CEFURONIUM (SODIUM) 250MG | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, meningococcal and other susceptible bacteria. | Prescription | | |

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|-----|--|---|---|----------------|----|--------|
| 6 | CEFURO INJECTION 750MG | CEFUROXIME (SODIUM) 750MG | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, meningococcal and other susceptible bacteria. | Prescription | | |
| 7 | CETADIME FOR INJECTION 2g | CEFTAZIDIME 2G | Ceftazidime is a bactericidal cephalosporin antibiotic that is resistant to a variety of endolines and is effective against a wide range of Gram-positive bacteria and Gram-negative bacteria. | Prescription | | |
| 8 | CETAXIME FOR INJECTION | CEFOTAXIME (SODIUM) 0.25GM/0.5GM/1GM/2 GM | Infections caused by Staphylococcus, Streptococcus, Streptococcus pneumoniae, meningococcal and other susceptible bacteria | Prescription | | |
| 9 | YUPIME POWDER FOR INJECTION | CEFEPIUME 0.5G/1G/2G | Inflammation caused by bacteria that are susceptible to Cefepime. | Prescription | | |
| 10 | CEFEPIUME POWDER FOR INJECTION "O-S" | CEFEPIUME 0.5G/1G/2G | Inflammation caused by bacteria that are susceptible to Cefepime. | Prescription | | |
| 11 | CEFZONE FOR IM/IV POWDER FOR INJECTION | CEFOPERAZONE 0.5G /1G | Treatment for upper and lower respiratory tract infection, upper and lower urinary tract infections, peritonitis, cholecystitis and other intraperitoneal infections, septicemia, meningitis, skin and soft tissue infections, bone, joint infections, pelvic inflammation, gonorrhea, endometritis and other genital tract infection | Prescription | | |

CARBAPENEM

| | | | | | | |
|---|---|-----------------------------------|---|--------------|--|--|
| 1 | MEOPEN POWDER FOR INJECTION 0.25G/ 0.5G/ 1G | MEROPENEM 0.25G/0.5G/1G | Inflammation caused by bacteria susceptible to Meropenem. | Prescription | | |
| 2 | YOUNAM POWDER FOR IV INJECTION | IMIPENEM 500MG + CILASTATIN 500MG | It has activity against inflammation of Gram-positive and Gram-negative organisms that are susceptible to imipenem. | Prescription | | |

TETRACYCLINES

| | | | | | | |
|---|------------------|-------------------------|---|--------------|--|--|
| 1 | DOMYCIN CAPSULES | DOXYCYCLINE (HCL) 115MG | The treatment of a variety of infections caused by susceptible strains of Gram-positive and Gram-negative bacteria and certain other micro-organisms. | prescription | | |
|---|------------------|-------------------------|---|--------------|--|--|

MACROLIDES

| | | | | | | |
|---|--------------------------------------|-------------------------------|--|--------------|--|--|
| 1 | ERYTH F.C. TABLET 500 MG "LONG LIFE" | ERYTHROMYCIN (ESTOLATE) 500MG | Infections caused by Streptococcus, Staphylococcus, Streptococcus pneumoniae, Pseudomonas aeruginosa, Treponema pallidum, Diphtheria bacillus, Listeria monocytogenes, Bacillus licheniformis, Legionella pneumophila, trachoma. | prescription | | |
|---|--------------------------------------|-------------------------------|--|--------------|--|--|

AMINO GLYCOSIDE

| | | | | | | |
|---|-------------------------|-----------------|---|--------------|--|--|
| 1 | GENTA 40MG/ML INJECTION | GENTAMICIN 40MG | The treatment of a variety of infections caused by susceptible strains of Gram-positive and Gram-negative bacteria and certain other micro-organisms. | prescription | | |
|---|-------------------------|-----------------|---|--------------|--|--|

POLYPEPTIDES

| | | | | | | |
|---|-----------------------------------|----------------------------|--|--------------|--|--|
| 1 | TECO LYOPHILIZED INJECTION | TEICOPLANIN 200 MG | It is an antibiotic that acts against coagulase producing Gram-positive cocci. It is used in the treatment of endocarditis, peritonitis, septicemia, osteoarthritis. | prescription | | |
| 2 | TEICOIN POWDER FOR I.V. INJECTION | TEICOPLANIN 200 MG 、 400MG | It is an antibiotic that acts against coagulase producing Gram-positive cocci. It is used in the treatment of endocarditis, peritonitis, septicemia, osteoarthritis. | prescription | | |

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|----------------------------------|--|--|---|----------------|----|-------------------|
| 3 | COLISTAR LYO INJ. | Colsitin Lyo inj. 2MIU | Used as a last-resort treatment for multidrug-resistant Gram negative infections including pneumonia | prescription | | |
| QUINOLONE | | | | | | |
| 1 | MOXACIN INFUSION SOLUTION 400MG/250ML | Moxifloxacin 400MG | Treatment for adults(>18yrs) inflammation caused by susceptible bacteria that includes 1.Upper and lower respiratory tract infection, such as sinusitis, acute exacerbation of chronic bronchitis, and Community-acquired pneumonia. 2.Infections of the skin and soft tissue. 3.Complicated intra-abdominal infection that includes many types of bacterial infections. | Prescription | | |
| 2 | MOLA INFUSION SOLUTION 400MG/250ML | MOXIFLOXACIN 1.6MG | Treatment for adults(>18yrs) inflammation caused by susceptible bacteria that includes, 1.Upper and lower respiratory tract infection such as, Sinusitis, acute exacerbation of chronic bronchitis, and Community-acquired pneumonia 2.Infections of the skin and soft tissue 3.Complicated intra-abdominal infection that includes many types of bacterial infections. | Prescription | | |
| 3 | LEVOFOR IV SOLUTION FOR INFUSION 5MG/ML (50ML/100ML/150ML) | LEVOFLOXACIN HEMIHYDRATE 5MG/ML (50ML/100ML/150ML) | Treat bacterial infections of 1.the lungs, in people with long-term breathing problems or Community-acquired pneumonia 2. urinary tract, including your kidneys or bladder 3. prostate gland, where you have a long lasting infection 4. skin and soft tissue, including muscles. | Prescription | | |
| 4 | OFLOKASIN 400 MG/250 ML SOLUTION FOR INFUSION | MOXIFLOXACIN 1.6 MG/ML 250 ML | It is used to treat infections caused by susceptible bacteria in adults (over 18 years old), including: upper and lower respiratory tract infections (acute sinusitis, acute exacerbation of chronic bronchitis, community pneumonia), skin and soft tissue infections, and complex intra-abdominal infections (including multiple bacterial infections). | prescription | | |
| MISCELLANEOUS ANTIBIOTICS | | | | | | |
| 1 | DAPTOMYCIN LYO INJ. 500MG | Daptomycin Lyo Inj. 500mg | Daptomycin is indicated for the treatment of the following infections. - Adult and paediatric (2 to 17 years of age) patients with complicated skin and soft-tissue infections (cSSTI). - Adult patients with right-sided infective endocarditis (RIE) due to <i>Staphylococcus aureus</i> . It is recommended that the decision to use daptomycin should take into account the antibacterial susceptibility of the organism and should be based on expert advice. Adult and paediatric (1 to 17 years of age) patients with <i>Staphylococcus aureus</i> bacteraemia (SAB) In adults, use in bacteraemia should be associated with RIE or with cSSTI, while in paediatric patients, use in bacteraemia should be associated with cSSTI. | prescription | | Under development |
| 2 | FOSMYCIN INJECTION | FOSFOMYCIN (SODIUM) 1G 2G 4G | Inflammation such as, Sepsis, bronchiolitis, Bronchiectasis, Pneumonia, lung abscess, Empyema, Peritonitis, Pyelonephritis, Cystitis caused by susceptible strains including <i>Pseudomonas aeruginosa</i> , <i>Proteus</i> , <i>Serratia</i> , <i>Staphylococcus</i> , and <i>E. Coli</i> . | prescription | | |

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|-----|---|------------------------------|--|----------------|----|-------------------|
| 3 | FOSFOMYCIN TROMETAMOL 3G GRANULES FOR ORAL SOLUTION | FOSFOMYCIN TROMETAMOL 3G | Ceftriaxone for Injection and Dextrose Injection is a cephalosporin antibacterial indicated for the treatment of the following infections caused by susceptible isolates of the designated bacteria: Lower Respiratory Tract Infections; Skin and Skin Structure Infections; Complicated and Uncomplicated Urinary Tract Infections; Pelvic Inflammatory Disease; Bacterial Septicemia; Bone and Joint Infections; Intra-abdominal Infections; Meningitis; and Surgical Prophylaxis. | prescription | | Under development |
| 4 | VANLYO POWDER FOR INJ. 0.5G 、 1G | VANCOMYCIN 0.5G 、 1G | Pseudomembranous colitis caused by endocarditis, osteomyelitis, pneumonia, septicemia, soft tissue infection, enteritis, and Clostridium infection caused by staphylococcal infection. | prescription | | |
| 5 | NUVANCO LYO INJ. 500MG/1G | Vancomycin Lyo Inj 500mg /1g | Endocarditis, osteomyelitis, pneumonia, septicemia, soft tissue infections, and enteritis caused by Staphylococcus infection; pseudomembranous colitis caused by Clostridium infection. | prescription | | |
| 6 | WINTORIN TABLETS | NALIDIXIC ACID 250MG | For the treatment of the urinary tract includes urethritis, cystitis, pyelitis, enteral infection, and biliary tract infection includes bacillary dysentery, enteritis, cholecystitis, cholangitis that caused by Gram-negative organisms. | prescription | | |

ANTIFUNGAL

| | | | | | | |
|---|---------------------------------------|--|---|--------------|--|--|
| 1 | CASPOFUNGIN/ANFARM for INJECTION 50MG | Caspofungin/Anfarm 50mg powder for concentrate for solution for infusion | <ul style="list-style-type: none"> •Treatment of invasive candidiasis in adult or paediatric patients. •Treatment of invasive aspergillosis in adult or paediatric patients who are refractory to or intolerant of amphotericin B, lipid formulations of amphotericin B and/or itraconazole. Refractoriness is defined as progression of infection or failure to improve after a minimum of 7 days of prior therapeutic doses of effective antifungal therapy. •Empirical therapy for presumed fungal infections (such as Candida or Aspergillus) in febrile, neutropaenic adult or paediatric patients. | prescription | | |
|---|---------------------------------------|--|---|--------------|--|--|

ANTICOAGULANTS

| | | | | | | |
|---|-------------------------------------|------------------------------------|---|--------------|--|-------------------|
| 1 | RIVAROXABAN F.C.TAB. 10MG/15MG/20MG | Rivaroxaban F.C.Tab 10mg/15mg/20mg | Indicated for prophylaxis of deep vein thrombosis (DVT), which may lead to pulmonary embolism (PE) in patients undergoing knee or hip replacement surgery. Indicated for prophylaxis of venous thromboembolism (VTE) and VTE related death during hospitalization and at discharge in adults admitted for an acute medical illness who are at risk for thromboembolic complications due to moderate or severe restricted mobility and other risk factors for VTE and not at high risk of bleeding. Indicated for treatment of existing DVT or PE. | prescription | | Under development |
|---|-------------------------------------|------------------------------------|---|--------------|--|-------------------|

ANTIDOTES

| | | | | | | |
|---|---------------------------------|--|--|--------------|----|-------------------|
| 1 | DEFERASIROX 125MG TAB. | Deferasirox 125mg Tab. | Treatment of chronic iron overload in patients 10 years of age and older with non-transfusion-dependent thalassemia (NTDT) syndromes and with a liver iron concentration (LIC) of at least 5 milligrams of iron per gram of liver dry weight (mg Fe/g dw) and a serum ferritin greater than 300 mcg/L. | prescription | BE | Under development |
| 2 | FOCAS LYO INJ. 50MG/100MG/150MG | Calcium Folinate 50mg 、 100mg 、 150mg Lyo Inj. | Treat or prevent serious blood cell disorders (such as thrombocytopenia, neutropenia, anemia) caused by certain drugs (folic acid antagonists such as methotrexate, trimethoprim, pyrimethamine). | prescription | | |

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|-----|-------------------|---------------------------|---|----------------|----|--------|
| 3 | CAFONIA TAB. 15MG | Calcium Folinate Tab.15mg | Calcium Folinate the immediate toxic effects of folic acid antagonists, e.g. Methotrexate. Calcium Folinate Rescue - a treatment technique using Calcium Folinate in conjunction with folic acid antagonists, e.g. methotrexate, to minimise systemic toxicity. The treatment of megaloblastic anaemias due to sprue, nutritional deficiency, pregnancy, infancy, liver disease and malabsorption syndrome. | prescription | | |

PSYCHIATRIC DRUG

| | | | | | | |
|----|---------------------------|---|--|--------------|----|-------------------|
| 1 | VENFAXIME S R.CAP. 75mg | VENLAFAXINE 75MG | Depression, pan-anxiety, and social anxiety. | prescription | BE | |
| 2 | PENTIXOL F.C. TABLETS | FLUPENTIXOL 3MG | Mental disorder | prescription | | |
| 3 | FLUXEL F.C. TABLETS | FLUPENTIXOL 0.5MG + MELITRACEN 10MG | Anxiety, depression. | prescription | | |
| 4 | ABIZOLE | Aripiprazole F.C.Tab. 10mg | Aripiprazole for the treatment of bipolar disorder (mania and mixed episodes and as maintenance treatment), as adjunctive treatment for major depressive disorder (2006) and for autism spectrum disorders | prescription | BE | |
| 5 | OLZAPINE F.C. TAB 10MG | OLANZAPINE 10MG | Schizophrenia and other apparently positive and / or negative psychosis, mania of bipolar disorder, prevention of bipolar disorder recurrence. | prescription | BE | |
| 6 | OLZAPINE F.C. TAB 5MG | OLANZAPINE 5MG | Schizophrenia and other apparently positive and / or negative psychosis, mania of bipolar disorder, prevention of bipolar disorder recurrence. | prescription | BE | |
| 7 | QUETIPINE F.C. TAB "L.L." | QUETIAPINE 200MG | Sympathetic disorders, episodes of bipolar disorder. | prescription | BE | |
| 8 | QUETIPINE | QUETIAPINE 100MG/25MG | Sympathetic disorders, episodes of bipolar disorder. | prescription | BE | Under development |
| 9 | SENTRA F.C. TAB. 100mg | SERTRALINE 100MG | Depression, obsessive-compulsive disorder, panic disorder, post-traumatic stress syndrome (PTSD), social phobia and premenstrual disease (PMDD Premenstrual Dysphoric Disorder. | prescription | BE | |
| 10 | SERULINE F.C.TAB 50MG | SERTRALINE 50MG | Depression, obsessive-compulsive disorder, panic disorder, post-traumatic stress syndrome (PTSD), social phobia and premenstrual disease (PMDD Premenstrual Dysphoric Disorder. | prescription | BE | |
| 11 | E-MINE TABLETS | NITRAZEPAM 10MG | Used for the short term treatment of sleeplessness | prescription | | |
| 12 | LUMRIN F.C. TABLETS | CHLORDIAZEPOXIDE HCL 10MG | Used for the short-term relief of anxiety, symptoms of alcohol withdrawal. | prescription | | |
| 13 | STIMIN F C TAB.10MG | ZOLPIDEM HEMITARTRATE 10MG | Insomnia. | prescription | BE | |
| 14 | SILTIN TABLETS 5MG | SELEGILINE HYDROCHLORIDE 5MG (DEPRENLY HYDROCHLORIDE) | Treatment of the early stages of Parkinson's disease. | prescription | | |

DEMENTIA THERAPEUTICS AGENTS

| | | | | | | |
|---|--------------------------------|---------------------------------|---|--------------|--|-------------------|
| 1 | MEMANTINE ORAL SOLUTION 2MG/ML | Memantine oral solution 2mg /ml | Memantine is used to treat moderate to severe confusion (dementia) related to Alzheimer's disease | prescription | | Under development |
|---|--------------------------------|---------------------------------|---|--------------|--|-------------------|

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|---------------------------------------|---|--|--|----------------|----|-------------------|
| FOR CARDIOVASCULAR DISEASE | | | | | | |
| 1 | ALADDIN F.C. Tablets 10mg | ROSVUVESTATIN CALCIUM 10MG | Treatment for hypercholesterolemia, hypertriglyceridemia . | prescription | BE | |
| 2 | TICLOUD F.C. TABLETS 100MG | TICLOPIDINE 100MG | Reduce the risk of thrombotic stroke (fatal or nonfatal) in patients who have experienced stroke precursors, and in patients who have had a completed thrombotic stroke. TICLOUD should be reserved for patients who are intolerant or allergic to aspirin therapy or who have failed aspirin therapy. | prescription | | |
| 3 | LIPOSEC CAPSULES 300MG (ETOFIBRATE) | ETOFLIBRATE 300MG | For hyperlipidemia | prescription | | |
| FOR GASTROINTESTINAL DISORDERS | | | | | | |
| 1 | LASUS ORAL SALINE LAXATIVE | MONOBASIC SODIUM PHOSPHATE 480MG + DIBASIC SODIUM PHOSPHATE 180MG | Relieve occasional constipation or bowel irrigation before rectal examination. | prescription | | |
| 2 | ASTREX CAP | CETRAXATE HCL 200MG | Cetraate prevents gastric mucosal blood flow decrease in H. pylori-infected patients. It is usually used to improve gastric mucosal lesion in acute gastritis or acute exacerbation of chronic gastritis and to treat gastric ulcer. | prescription | | |
| 3 | METHYLSkopola MINE METHYLSULFATE F.C. TABLETS "L.L." | METHSCOPOLAMINE METHYLSULFATE 1MG | Treatment for gastritis, gastric ulcer, peptic ulcer, cramps and pain caused by above symptoms. | prescription | | |
| 4 | KORINTI TABLETS | DICYCLOMINE HCL 2.5MG MAGNESIUM OXIDE 100MG ALUMINUM HYDROXIDE GEL 200MG | Treatment for gastric ulcer, peptic ulcer, hyperacidity, ulcerative colitis, spastic colitis. | OTC | | |
| 5 | EMOSE CAPSULES 2MG (LOPERAMIDE) | LOPERAMIDE HCL 2MG | Relieve the symptoms of sudden, short term diarrhea ("acute" diarrhea) | prescription | | |
| 6 | SUWEYLIN TABLETS | METHYLMETHIONINE SULFONIUM CYCLAMATE 50MG BELLADONNA EXTRACT 5MG DRIED ALUMINUM HYDROXIDE GEL 40MG CALCIUM CARBONATE 110MG CHLOROPHYLLIN 15MG | Treatment for acute gastritis, chronic gastritis, gastric ulcer, peptic ulcer and hyperacidity | OTC | | |
| 7 | SULSON S.C. TABLETS 100MG | MEBEVERINE HCL 100MG | Treatment for the symptoms of gastrospasm, enterospasm, chronic irritable colon, gastritis, gastric ulcer, peptic ulcer, mucous colitis, spastic colitis, biliary dyskinesia. | prescription | | |
| 8 | ESOCOMFORT E.F.C TAB 40MG | ESOMEPRAZOLE 40MG | Esomeprazole magnesium oral capsule is used to treat conditions caused by too much acid production in your stomach. These conditions include gastroesophageal reflux disease (GERD), erosive esophagitis, Zollinger-Ellison syndrome, and Helicobacter pylori infections. | prescription | BE | |
| 9 | ESOCOMFORT E.F.C TAB 20MG | ESOMEPRAZOLE 20MG | Esomeprazole magnesium oral capsule is used to treat conditions caused by too much acid production in your stomach. These conditions include gastroesophageal reflux disease (GERD), erosive esophagitis, Zollinger-Ellison syndrome, and Helicobacter pylori infections. | OTC | BE | Under development |
| 10 | OMEPRAZOLE LYO INJ. 40MG | Omeprazole LYO INJ 40MG (10cc vial) | This medicine is used to treat and help heal peptic ulcers. Used to help prevent gastric or duodenal ulcers from coming back. To treat a rare condition called Zollinger-Ellison syndrome | prescription | | |

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|--|--|-----------------------------------|---|----------------|----|--------|
| 11 | PANTOPRAZOLE LYO INJ. 40MG | Pantoprazole Lyo Inj.40mg | Pantoprazole injection is used as a short term treatment to treat gastroesophageal reflux disease (GERD; a condition in which backward flow of acid from the stomach causes heartburn and possible injury of the esophagus [the tube between the throat and stomach]) in people who have had damage to their esophagus and who are unable to take pantoprazole by mouth. It is also used to treat conditions where the stomach produces too much acid, such as Zollinger-Ellison syndrome | prescription | | |
| FOR LIVER, GALLBLADDER, AND HEPATITIS | | | | | | |
| 1 | ENITOL F.C. TABLETS | ANETHOLE TRITHIONE 12.5MG | The treatment of cholecystitis, gallstones, biliary dyskinesia. | prescription | | |
| 2 | UROSE CAPSULES 300MG | URSODEOXYCHOLIC ACID 300MG | To dissolve certain types of gall stones (stones in the gall bladder), which contain cholesterol, and treat a type of liver disease known as primary biliary cirrhosis(PBC). | prescription | | |
| 3 | BARAVIR F.C. Tablets 0.5MG | ENTECAVIR 0.5MG | Treat chronic (long term) hepatitis B virus (HBV) infection in adults and children aged more than 2 years | prescription | BE | |
| 4 | BARAVIR F.C. Tablets 1MG | ENTECAVIR 1MG | Treatment of adults with signs of hepatitis B virus replication and chronic hepatitis B patients over 2 years of age. | prescription | BE | |
| LIVER PROTECTANT | | | | | | |
| 1 | DATHIONE LYO INJ. 500MG | Glutathione Reduced Lyo Inj.500mg | Glutathione injections is anti-aging injections to help fight the signs of aging. It can also help reverse the signs of free radicals in the body and prevent oxidative stress. | prescription | | |
| FOR DIABETES | | | | | | |
| 1 | GLAZIDE TABLETS 80MG (GLICLAZIDE) | GLICLAZIDE 80MG | Diabetes | prescription | | |
| FOR GOUT | | | | | | |
| 1 | BENMARONE TABLETS 50MG | BENZBROMARONE 50MG | Treatment for gout and hypertension caused by hyperuricemia | prescription | | |
| MUSCLE RELAXANT | | | | | | |
| 1 | MAOLUX TABLETS 125MG (CHLORPHENESIN) | CHLORPHENESIN CARBAMATE 125MG | Spasm caused by locomotive organs such as, low back pain, spondylosis deformans, osteoporosis of spine | prescription | | |
| ANTIHISTAMINE | | | | | | |
| 1 | BROMPHENIRAMINE MALEATE TABLETS 4MG "L.L." | BROMPHENIRAMINE MALEATE 4MG | Relieve symptoms of allergy, and hay fever. These symptoms include runny nose, sneezing, itchy eyes/throat, and itchy skin caused by allergy, rash. | OTC | | |
| 2 | CHLORPHENIRAMINE MALEATE TABLETS "S.S.P." | CHLORPHENIRAMINE MALEATE 4MG | Relieve symptoms of allergy, and hay fever. These symptoms include runny nose, sneezing, itchy eyes/throat, and itchy skin caused by allergy, rash. | OTC | | |
| IMMUNOSUPPRESSION | | | | | | |
| 1 | AZAPRINE F.C. Tablets 50mg | AZATHIOPRINE 50MG | Help your body accept kidney transplant, treat systemic lupus erythematosus, rheumatoid arthritis, acute leukemia, and chronic leukemia. | prescription | | |

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|-----|-------------------------------|-----------------------|--|----------------|----|--------|
| 2 | Cycos SOFT GELATIN CAP. 100MG | CYCLOSPORIN 100MG | Prevention of transplant rejection after organ transplantation and bone marrow transplantation, prevention of graft-versus-host disease, activity of blindness, middle or posterior non-infectious uveitis, uncontrolled by traditional therapy, repeated inflammation of BEHCT disease, and invasion of retina Severe dryness that is ineffective or unsuitable for alternative therapy; severe rheumatoid arthritis that is ineffective or unsuitable for standard therapy; primary renal syndrome with steroid-ineffective or steroid-dependent (Biopsy) A small lesion disease or localized glomerulosclerosis, a patient who is ineffective by Cytostatic therapy and has a renal function index above 50% of normal. | prescription | BE | |

FOR FLU

| | | | | | | |
|---|----------------------------|---|--|--------------|--|-------------------|
| 1 | KAN MAW KEH TABLETS "L.L." | ACETAMINOPHEN 100MG ETHOXYBENZAMIDE 83.33MG NOSCAPINE 5MG DL-METHYLEPHEDRINE HCL 5MG CAFFEINE ANHYDROUS 15MG CHLORPHENIRAMINE MALEATE 1.25MG | For the temporary relief of minor aches and pains associated with runny nose, stuffy nose, sneeze, cough, mucus, sore throat, fever, headache, arthritis pain, muscular aches. | OTC | | |
| 2 | ASPHONLIN CAPSULES | ACETAMINOPHEN 500MG DEXCHLORPHENIRAMINE MALEATE 2MG PSEUDOEPHEDRINE HCL 60MG DEXTROMETHORPHAN HBR 15MG | For the temporary relief of minor aches and pains associated with stuffy nose, runny nose, sneeze, sore throat, cough, fever, arthritis pain, muscular aches. | OTC | | |
| 3 | COMCOD SOLUTION | ACETAMINOPHEN 15MG CAFFEINE ANHYDROUS 1.24MG CHLORPHENIRAMINE MALEATE 0.124MG DL-METHYLEPHEDRINE HCL 1MG | For the temporary relief of minor aches and pains associated with sore throat, fever, headache, arthritis pain, muscular aches, stuffy nose, runny nose, sneeze. | OTC | | |
| 4 | OSLETAMIVIR 75MG CAP | Osletamivir 75mg Cap | Treatment of acute, uncomplicated illness due to influenza A and B infection in patients 2 weeks of age and older who have been symptomatic for no more than 48 hours. | prescription | | Under development |
| 5 | ANESTAT CAPSULES | ACETAMINOPHEN 320MG CAFFEINE ANHYDROUS 40MG SALICYLAMIDE 200MG | Fever reducer and pain reliever includes headache, toothaches, sore throat, arthritis, muscular aches, and dysmenorrhea. | OTC | | |

ENT Drugs

| | | | | | | |
|---|------------------------------|------------------------------|--|--------------|--|-------------------|
| 1 | POVIDONE LODINE THROAT SPRAY | Povidone Iodine Throat Spray | Indicated for the treatment of acute mucosal infections of the mouth and pharynx including stomatitis, gingivitis, aphthous ulcers, pharyngitis, tonsillitis, monilial infections, common colds and influenza. | prescription | | Under development |
|---|------------------------------|------------------------------|--|--------------|--|-------------------|

BRONCHODILATORS AND ANTIASTHMATIC DRUGS

| | | | | | | |
|---|---------------------------------|--------------------------|---|--------------|----|--|
| 1 | MONAST FILM-COATED TABLETS 10MG | Montelukast (Sodium)10mg | Montelukast is used to control and prevent symptoms caused by asthma (such as wheezing and shortness of breath). It is also used before exercise to prevent breathing problems during exercise (bronchospasm). This medication can help decrease the number of times you need to use your quick relief inhaler. | prescription | BE | |
|---|---------------------------------|--------------------------|---|--------------|----|--|

URINARY SYSTEM

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|---|----------------------------------|---|--|----------------|----|-------------------|
| 1 | DUTASTERIDE SOFT CAP 0.5MG | Dutasteride Soft Cap 0.5mg | indicated for the treatment of symptomatic benign prostatic hyperplasia (BPH) in men with an enlarged prostate to: improve symptoms, reduce the risk of acute urinary retention (AUR), and. reduce the risk of the need for BPH-related surgery. | prescription | BE | Under development |
| 2 | Xinlis film-coated tablets 20 mg | Tadalafil F.C.Tab. 20mg | Treat erection problems (erectile dysfunction) and symptoms of an enlarged prostate (benign prostate enlargement). | prescription | BE | |
| 3 | Xinlis film-coated tablets 5 mg | Tadalafil F.C.Tab. 5mg | Treat erection problems (erectile dysfunction) and symptoms of an enlarged prostate (benign prostate enlargement). | prescription | BE | |
| ARTERIAL INJECTION | | | | | | |
| 1 | ALPROSTADIL | Alprostadil 0.02mg Lyo | For temporary maintenance of ductus arteriosus patency until corrective or palliative surgery can be performed in neonates with ductal-dependent congenital heart defects. | prescription | | Under development |
| ANTITUSSIVE, MUCOLYTICS | | | | | | |
| 1 | NOSCAPINE TABLETS | NOSCAPINE 10MG | Relieving cough | OTC | | |
| 2 | SU CO TABLETS 20 MG | EPRAZINONE HCL 20MG | Relieving cough, thinning the mucus | OTC | | |
| 3 | BISUCON TABLETS 8MG | BROMHEXINE HCL 8MG | Thinning the mucus in the airways, thus making the mucus less sticky and it also facilitates the removal of the mucus. | OTC | | |
| PAIN RELIEVER, ANTI-INFLAMMATORY | | | | | | |
| 1 | Xingaba Capsules 75mg "L.L." | PREGABALIN CAP. 75MG | It is used to help control certain kinds of seizures. It is used to treat painful nerve diseases. It is used to treat fibromyalgia. | prescription | BE | |
| 2 | DICLOFENAC TABLETS 25MG "L.L." | DICLOFENAC SODIUM 25MG | Relieve pain, reduce swelling and ease inflammation in conditions affecting the joints, muscles and tendons. | prescription | | |
| FOR THROAT AND MOUTH | | | | | | |
| 1 | DROCHEN LOZENGES 0.5MG | DOMIPHEN BROMIDE 0.5MG | Treatment of sore throat caused by Stomatitis, inflammation of throat, and thicken mucus. | OTC | | |
| STEROID | | | | | | |
| 1 | BETASHIN TABLETS | BETAMETHASONE 0.5MG LACTOSE 75MG STARCH 51.5MG MAGNESIUM STEARATE 3MG | Treatment for rheumatoid arthritis, allergic reactions, connective tissue disorders, inflammatory conditions of the skin, inflammatory conditions of the eye, and asthma. | prescription | | |
| ANTHELMINTICS | | | | | | |
| 1 | POKIUU TABLETS | PYRVINIUM PAMOATE 75MG | The treatment of enterobiasis caused by Enterobius vermicularis | OTC | | |
| ORALBASE, AND SKIN OINTMENT | | | | | | |
| 1 | ORALFIX ORAL BASE | TRIAMCINOLONE 1MG | Gingivitis, oral mucosa, tongue inflammation or ulcers, angular cheilitis, pyorrhea | OTC | | |
| 2 | DOXEFEN CREAM 50MG/GM | DOXE PIN HCL 50MG | Relieve itching associated with eczematous dermatitis. | prescription | | |
| 3 | TONIN GEL 100MG/GM (ETOGENAMATE) | ETOGENAMATE 100MG | Relieve rheumatoid arthritis, upper wrist scapular inflammation, low back pain, sciatica, tenosynovitis, and bursitis. | prescription | | |

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|-----|-------------------------|--|--|----------------|----|-------------------|
| 4 | MOMO CREAM | MOMETASONE FUROATE 1MG/G | Reduce redness and itchiness caused by certain skin problems that susceptible of the steroid. | prescription | | |
| 5 | NAFINE CREAM | NAFTIFINE 10MG/G | 1.Infections caused by trichophyton species, small spore ringworm species, hair flocculent epidermoid fungi in the skin and its affiliated tissue. 2.Epidermal candidiasis 3.Mycomorphosis 4.Pityriasis versicolor | prescription | | |
| 6 | DERMALIN CREAM | BETAMERTHASONE DIPROPIONATE 0.64MG (eq to BETAMETHASONE 0.5MG)+CLOTRIMAZOLE 10MG | Indicated for the topical treatment of symptomatic inflammatory tinea pedis, tinea cruris, and tinea corporis due to Epidermophyton floccosum, Trichophyton mentagrophytes and Trichophyton rubrum. | prescription | | |
| 7 | TROSTEN CREAM 1% "L.L." | TCOCONAZOLE 10MG | Treat infections caused by certain fungi. | prescription | | |
| 8 | SEMPARIN OINTMENT | TOLNAFTATE 20MG CHLORHEXIDINE HCL 2MG | Topical antifungal for the treatment of Athlete's Foot, tinea cruris, tinea versicolor. | OTC | | |
| 9 | SSD CREAM " DI" | SILVER SULFADIAZINE 10MG/G | Prevent infection and treatment in burns, especially for Gram-negative bacterial infections, such as pseudomonas aeruginosa infection. Also for stab wounds, trauma, and other types of wounds. | prescription | | |
| 10 | ACNEAN CREAM 200MG/G | AZELAIC ACID 200MG | Acne vulgaris | prescription | | |
| 11 | B.P. GEL 10% | BENZOYL PEROXIDE 100MG | Treatment of acne vulgaris | OTC | | |
| 12 | SHEATE GEL 5% | BENZOYL PEROXIDE 50MG | Treat moderate acne (spots). | OTC | | |
| 13 | EUCLOBE CREAM 0.5MG/GM | CLOBETASOL-17-PROPIONATE 0.5MG | Treatment of certain skin problems include eczema, psoriasis and discoid lupus erythematosus. | prescription | | |
| 14 | FLUSONE CREAM 0.05% | FLUTICASONE PROPIONATE 0.5MG | Relieve inflammation and itching symptoms of skin diseases, such as eczema, including atopic dermatitis, early stage of eczema and discoid eczema, nodular prurigo, psoriasis which is not including plaque psoriasis, neurodermatitis includes Lichen planus, liposomal dermatitis, contact allergic reactions, discoid erythema lupus erythematosus, dermatitis that has not responded to milder steroid creams, insect bites, and prickly heat. | prescription | | |
| 15 | BUTEFINE CREAM 10MG/G | BUTENAFINE HYDROCHLORIDE 10MG | The topical treatment of the following superficial dermatophytosis: interdigital tinea pedis (athlete's foot), tinea corporis (ringworm) and tinea cruris (jock itch) due to E. floccosum, T. mentagrophytes, T. rubrum, and T. tonsurans. | prescription | | |
| 16 | MUMBAI DEPIGMENT CREAM | DEXAMETHASONE 0.3MG/ G HYDROQUINONE 50MG, RETINOIC ACID 0.3MG, | Indicated for the gradual bleaching of hyperpigmented skin conditions such as berloque dermatitis, contact dermatitis, atopic dermatitis, freckles, seborrheic keratosis chloasma, riehls melanosis, and other unwanted areas of melanin hyperpigmentation. | prescription | | |
| 17 | PERMETHRIN 5% | Permethrin 5% x 30g | Permethrin 5% w/w Cream is indicated for the treatment of scabies. | prescription | | under development |
| 18 | ACURE GEL 0.1% | ADAPALENE 0.1% | TRETMENT FOR ACNE | prescription | | |

FOR CANCER

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|------------|--|---|---|----------------|--------------------|--------|
| 1 | DOXORUBICIN HCL 10MG | Doxorubicin Hcl 10mg | <ul style="list-style-type: none"> Doxirubicin Hcl is the treatment of AIDS related Kaposi's sarcoma patients with low CD4 (<200 CD4, lymphocytes/mm3) and mucosal, skin or internal organs. Treatment for Ovarian cancer, after failure of platinum-based chemotherapy. Doxorubicin can be used as first line treatment of patients with metastatic breast cancer who are at risk of heart disease. Doxorubicin, in combination with bortezomib is indicate multiple myeloma. | prescription | | |
| 2 | EPIRUBICIN HCL 1MG/10MG/50MG | Epirubicin Hcl 1mg/ 10mg/ 50mg | <ul style="list-style-type: none"> Epirubicin Hydrochloride has produced responses in a wide range of neoplastic conditions including breast, ovarian, gastric, lung and colorectal carcinomas, malignant lymphomas and multiple myeloma. | prescription | | |
| 3 | OXAOXA INJ 5mg/ml | Oxaliplatin 5mg/ml | <p>Oxaliplatin in combination with 5-fluorouracil (5-FU) and folinic acid (FA) is indicated for:</p> <ul style="list-style-type: none"> adjuvant treatment of stage III (Dukes' C) colon cancer after complete resection of primary tumour. treatment of metastatic colorectal cancer. | prescription | | |
| 4 | PEMETREXED DISODIUM HEPTAHYD 100MG/500MG | Pemetrexed Disodium Heptahyd 100mg/ 500mg | <ul style="list-style-type: none"> Pemetrexed Disodium Heptahyd in combination with cisplatin is indicated for the first line treatment of patients with locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology. Pemetrexed Disodium Heptahyd is indicated as monotherapy for the maintenance treatment of locally advanced or metastatic non-small cell lungcancer other than predominantly squamous cell histology in patients whose disease has not progressed immediatelyfollowing platinum-based chemotherapy. Pemetrexed Disodium Heptahyd is indicated as monotherapy for the second line treatment of patients with locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology. Pemetrexed Disodium Heptahyd in combination with cisplatin is indicated for the treatment of chemotherapy naïve patients with unresectablemalignant pleural mesothelioma. | prescription | | |
| 5 | IMATIS F.C. TAB. 100MG | Imatinib 100mg Tab. | <p>(for simple indication)</p> <ul style="list-style-type: none"> treatment of patients with newly diagnosed Philadelphia chromosome positive chronic phase chronic myeloid leukemia (CML), treatment of patients with Philadelphia chromosome positive chronic myeloid leukemia (CML) in accelerated phase, treatment of patients with Philadelphia chromosome positive chronic myeloid leukemia (CML) in blast phase, treatment of patients with Philadelphia chromosome positive chronic myeloid leukemia (CML) who are resistant to other therapies, treatment of patients with chronic myeloid leukemia (CML) in chronic/accelerated/blast phase who were initially diagnosed with Philadelphia chromosome positive chronic myeloid leukemia (CML) that has, however, become Philadelphia chromosome negative due to the treatment, first-line treatment of children at and over 3 years of age with chronic myeloid leukemia (CML), the treatment of adult patients with Kit (CD 117) positive unresectable and/or metastatic malignant gastrointestinal | prescription | Under registration | |
| 1 | XINVIR LYO INJ. 250MG | Acyclovir Lyo Inj.25mg (10cc vial) | Indicated for the treatment of initial and recurrent mucosal and cutaneous herpes simplex (HSV-1 and HSV-2) in immunocompromised patients. Indicated for the treatment of herpes simplex encephalitis. Indicated for the treatment of neonatal herpes infections. | prescription | | |
| Hemorrhoid | | | | | | |

| No. | Product Name | Ingredients/Strengths | Indication | Classification | BE | Remark |
|--|-----------------------------------|-----------------------|--|----------------|----|-------------------|
| 1 | Aililon film-coated Tablets 500mg | Diosmin F.C.Tab.500mg | Helps improve local swelling or pain caused by chronic venous insufficiency and relieve hemorrhoid symptoms. | OTC | | Under Development |
| ANTI-INFLAMMATORY AND PAIN RELIEF PATCHES | | | | | | |
| 1 | LIDOCAINE PATCH 5% | LIDOCAINE 700MG/14G | Used to relieve post-herpetic neuralgia. The site of use is limited to undamaged skin. | prescription | | |

 LONG LIFE NUTRACEUTICAL INT'L CO., LTD.

<https://www.lehyinn.com/>

Contact us:

Address: No.119, Hwai An St., San Min Dist. Kaohsiung Taiwan 80762

Phone: +886 7 3863323

Fax: + 886 7 3867999

Email : service@lehyinn.com